




TRIM WITH BROMELAIN:

PINE BOTTLE

LIPOLYSIS REVOLUTION



Efficiency

Revitalization


Precision

Versatility

Comprehensive

TRIM WITH BROMELAIN:

Lipolysis Revolution, Pine Bottle



Enhance Skin Elasticity, Reduce Wrinkles.

Pine Bottle enhances skin elasticity and reduces wrinkles. Unlike typical lipolysis products that can cause wrinkles during fat dissolution, our solution, combined with Riboflavin, Cyanocobalamin, and L-Carnitine, accelerates lipolysis, safely dissolving fat cells and addressing potential wrinkles.

Fat Breakdown

Pine Bottle features potent ingredients like Lecithin, Bromelain, and Riboflavin (Vitamin B2) for effective fat breakdown. Lecithin's lipophilic tails attract and absorb fats, playing a key role, while its stimulation of lipase release aids in breaking down triglycerides into energy-providing fatty acids for the body's cells.

GradualismComfortCustomizationAdaptabilityInnovation

Specification

Product Name	PINE BOTTLE
Capacity	10cc/vial (5vials/box)
Main Ingredients	Lecithin, Bromelain, Riboflavin

Main Ingredients

Lecithin

Derived from soybeans and egg yolks, lecithin effectively breaks down excess body fat into energy-usable fatty acids, promoting fat burning and diminishing the risk of fat storage in adipose tissue.

Bromelain

Bromelain, anti-inflammatory and anti-edema effects. Bromelain is a mixture of enzymes found naturally in the juice and stems of the pineapple plant. It helps to break down proteins and reduce inflammation.

Riboflavin

Riboflavin (Vitamin B2) activates fat metabolism by serving as electron carriers in energy production, antioxidant function, and various metabolic pathways. Essential for cellular growth and function, riboflavin aids in converting nutrients into energy.

Lipolysis Revolution

PINEBOTTLE



Total Usage Volume

Treatment Area	Dosage of point	Depth	Interval of Point	Number of Point
Face (Sub-mental)	0.1~0.25cc (0.5~0.8cc)	2~4mm	0.5~1.0cm	20~25 points
Body	1.0~2.0cc	6~12mm	1.5~2.0cm	30~70 points


Procedure of Process



Treatments occur 1 to 3 times, with a 4-week gap between sessions. Post the 3rd treatment, the interval increases, and additional procedures are tailored to the patient's needs. Gradual adaptation is emphasized over rapid fat reduction.

Combination Therapy

Ultrasound softens fat cells, aiding breakdown and preventing clumping. Combined with cryo-fat decomposition and small-dose DC therapy, it offers unique benefits. Frozen state procedures minimize pain. To reduce pain, a thin needle(e.g. 34G 2mm needle) and alcohol ice gauze are used. Diluting 1-2cc of lidocaine per 10cc, along with less than 1cc of triamcinolone, significantly reduces pain and swelling.



Facial Treatment

The face is divided, treating left and right sides along the jawline. Using 0.1 to 0.25 ml per point, four lines are designed on each side, totaling approximately 5.0ml per side. The amount can be adjusted based on fat levels, using one vial for both procedures.

Total: 10.0ml is used

Forearm Treatment

The amount used is consistent with facial procedures, applying 0.5ml to 1.0ml per point. About 20 points on each forearm, including 10 on the inner and lower areas, are treated. For one side, approximately 15.0ml is used, with a total of 20.0~30.0ml for both forearms.

Total: 20.0~30.0ml used

Abdominal Treatment

Approximately 1.0 to 2.0ml per point is used, totaling about 50ml for the treatment. The upper abdomen receives approximately 20 points centered on the navel, with 15 points on each side and 10 points on each side area.

Total: 30.0~50.0ml used

Thigh Treatment

Similar to forearm treatment, the inner and lower thighs are treated. Administering approximately 10.0 to 20.0 ml per thigh, about 20 to 30 points are treated. Each thigh receives approximately 10.0ml, totaling 20.0ml for the entire procedure.

Total: 20.0~40.0ml used

Hip-up Treatment

The procedure, above the buttocks' lower line (banana line), uses 0.5 to 1.2ml at 1.5cm intervals along the line. Repeated on the opposite hip, using 10-15 points per hip, approximately 10ml is used. Pine Bottle removes fat from sagging buttock skin.

Total: 10.0~20.0ml used

FAQ

Frequently Asked Questions

Q1

What is Pine Bottle and How does it work?

Pine Bottle, a lipolysis solution with Riboflavin, Bromelain, and Lecithin, effectively reduces fat. Recommended for face and body application.

Q2

How does Pine Bottle differ from others?

Unlike formulas with potential side effects, Pine Bottle is free from DC/PPC contents. Manageable swelling and pain may occur without disrupting daily life.

Q3

Are there any benefits beyond Lipolysis?

Pine Bottle offers skin-lifting, wrinkle removal, and improved elasticity. Enhances overall skin health and brightness.

Q4

Are there any precautions during application?

Avoid Pine Bottle on inflamed areas. Not for use during pregnancy, severe diabetes, or poor health.

Q5

Post-Application Care?

Refrain from intense activities and sauna on the first day.
Avoid irritation in the treated area. Reduce facial swelling with gentle lymph massage.
Apply sunscreen outdoors.

TRIM WITH BROMELAIN:

PINE BOTTLE

Specification

Product Name	PINE BOTTLE
Capacity	10cc/vial (5vials/box)
Main Ingredients	Lecithin, Bromelain, Riboflavin

Caution for Use:

Storage:

- Room temperature, avoid sunlight.
- Refrigerate opened products; use within 3 days.

General Information:

- Avoid eye contact.
- Dispose after use; no reuse.
- Shake vial before use.
- 3 sessions, 4-6 week intervals.
- Immediate ampoule use; no mixing.
- Dosage: not over 3.0ml at a spot.

Absolute Contraindications:

- Anticoagulation, chemotherapy, pregnancy.
- Children under 18, severe chronic illness, soy allergy.
- Immune deficiency, open sores.

Relative Contraindications:

- Inability to maintain results.
- Previous surgery, extreme skin laxity.
- Type II diabetes, autoimmune disease.
- Inability to follow treatment plan

Cautions After Treatment:

- No tobacco, alcohol, spicy foods.
- Expect redness, swelling, bruising.
- Keep area clean; no water for 24 hours.
- Effect in 1-2 weeks; increased thirst/urination normal.
- Soy allergy caution; extreme allergy—use with care.
- Drink more water; avoid irritating skin procedures.
- Post-procedure, avoid friction, tight clothes, alcohol.
- No late-night meals; light exercise, massage recommended.
- Use hot towel/sauna for 2-3 days.
- Permanent effect with no weight gain; rebound possible without diet attention.

Expected Side Effects:

- Swelling, bruising for 10 days.
- Temporary worsening before improvement.
- Dose-related nausea, diarrhea, flu-like symptoms.
- Jiggly tissue for 3-5 days, subcutaneous nodules at 2 weeks.
- Temporary tissue stiffness; no improvement for 4-6 weeks.

Complications:

- True complications not expected.
- Hyperpigmentation, skin loss, visible blood vessels possible.
- Persistent pain, skin irregularity, less improvement.
- Persistent nodularity

TRIM WITH BROMELAIN:

PINE BOTTLE