PRIME
FAT DISSOLVING
SOLUTION
Prime
DR.LIPO+
2ml x 10 Vials

Series of **DR.LIPO+**

Always excellent and expected results

Designed to eliminate local fat deposits on the face and body.

Harmless Ingredients

Painless pH Level 7.0

Unique anti-puffiness formula

Why *Prime DR.LIPO+?*

- ✓ High efficiency in affected areas.
- ✓ Ability to work with any part of the body and face.
- ✓ Excellent tolerability due to the mild effect of the product components on tissues.
- ✓ Short rehabilitation period after product administration.
- ✓ Improved lymphatic drainage properties due to the use of muscle relaxants in the composition of the product.
- ✓ Ingredients of the highest quality and purity.
- ✓ During clinical trials, no side effects or complications were identified.

PRIME FAT DISSOLVING SOLUTION
PRIME FAT DISSOLVING SOLUTION
PRIME FAT DISSOLVING SOLUTION

How does *Prime DR.LIPO+ work?*

"By selecting balanced and complementary ingredients, Prime has a distinct advantage over competitive products."

Muscle relaxants reduce skin laxity, reduce wrinkles and improve lymphatic drainage. Cell growth factors promote increased tissue regeneration and initiate the processes of neocollagenesis. Antioxidants fight free radicals and promote tissue health, while vitamins help improve elasticity, improve skin turgor and rejuvenate the skin.




1
BEFORE THERAPY
Cumulative fat cells
2
1-5 DAYS
Primary injection into the fat layer
3
1 WEEK LATER
Lysis of fat cells
4
2 WEEKS LATER
Speeding up the lysis process
5
4 WEEKS
The natural removal of reduced fat cells
DIRECT ADIPOCYTE EFFECTS (UNTIL 2DAYS)
LOCAL TISSUE RESPONSE
MULTIPLE TRATMENTS

BEFORE

AFTER

Advantages of *Prime*




Aged skin


New collagen production


Activation of cell growth factors


Cell regeneration

High Purity Lipolysis

Recovery of collagen and elastin (growth factor)

Lipolysis promoter

Skin antioxidant

Increase elasticity & prevent wrinkles

Calming & anti-inflammatory effect

Treatment area



FACE AREA

- Forehead area
- Periorbital zone
- Tip of the nose (depth 4mm)
- Zygomatic zone
- Chin
- Lower third of the face

BODY AREA

- Armpit Forearm
- Abdomen

BACK SIDE

- Back
- Waist
- Hips Buttocks
- Fibrous cellulite
- Outer thighs Inner thighs

Instruction

PRE-TREATMENT INSTRUCTIONS

1. Avoid certain products 2 days before and after treatment.
2. Shave the treatment area one day before if applicable.
3. Do not apply creams, lotions, or makeup to the treatment areas.
4. Ensure a recent meal before the procedure to prevent lightheadedness.
5. Schedule appointments before special events for bruising consideration.
6. Reschedule for severe rashes, cold sores, or blemishes in the treatment area.
7. Avoid specific medications and alcohol before treatment.

WORK PROTOCOL



	FACE	BODY
Spread Infiltration	Evenly spread the injection	
Techniques of Introduction	Pinching technique for injections	
Needle Type	Needle (30Gx1inch), Cannula, Meso-Roller or RF Device	
Depth (Fat Layer)	0.5-1cm	0.5-2cm
Angle of Inclination	90° (vertical direction)	
Volume	0.2cc - 1cc/cm²	
Total Volume	5cc - 20cc/time	Up to 40cc
Interval	2-4 weeks	
Total Schedule	3-6 times	

STEP**01** PREPARE SKIN FOR THE PROCEDURE

1. Cleaning and Shaving:
   - Clean the treatment area with ethanol.
   - Shave the entire target area.
2. Design:
   Draw the divided region for 1 ampoule and draw a 1cm grid on the target area.
3. Anesthesia:
   Administer 2% Lidocaine injection for local anesthesia (optional).

STEP**02** ADMINISTRATION



1. Choose the appropriate injection method based on the treatment plan and patient needs. Options include spread by cannula, injection by needle, or application by meso-roller.
2. Insert the needle or cannula into the fat layer at a depth recommended for the specific area being treated. Maintain a consistent distance of 1cm to 2cm between injection points for even coverage.
3. Administer the lipolytic solution with volumes ranging from 0.2cc to 1cc per square centimeter. Adhere to the recommended total volume of 5cc to 20cc per treatment session (depending on the thickness of subcutaneous fat tissue).
4. Depending on the chosen method, employ specific injection techniques:
   - For needle injections: Use a pinching technique, insert a 1-inch 30G needle in a vertical direction, and perform spread infiltration.
   - For cannula spread: Insert the cannula with medicine and perform backward spreading.
   - For meso-roller application: Apply the meso-roller evenly across the treatment area.

STEP**03** POST-INJECTION CARE

1. Avoid strenuous exercise or straining for 48 hours.
2. Avoid medications and supplements that may increase bruising for 48 hours.
3. Apply ice for the first day to reduce discomfort and swelling.

LIMITATIONS OF USE

- The safe use of Prime DR.LIPO+ for subcutaneous fat outside the submental region is not recommended.
- Dosage and Administration (Prime DR.LIPO+)
- Follow the dosage and administration guidelines provided for Prime DR.LIPO+.

POST-TREATMENT INSTRUCTIONS

- Avoid extreme heat for 24 hours.
- Do not consume alcohol or sodium for 3 days.
- Avoid strenuous exercise for 24 hours.
- Sleep on your back with an elevated head for decreased swelling.

WHAT TO EXPECT

- Swelling and bruising for 3-5 days, decreasing over weeks.
- Multiple treatments may be needed for the desired results.
- Ice packs are recommended for the first 12 hours.
- Do not massage the injection site.
- Results are not immediate; full results may take up to 12 weeks.

Warning ⓘ

- Do not use this solution earlier than 2 weeks after the administration of other lipolytics.
- Avoid combining Prime DR.LIPO+ with botulinum toxin preparations, biorevitalizers or HA fillers.
- It is not recommended to use the bolus technique in the face, as this may cause an uneven effect of the product on the pancreas.
- It is not recommended to administer the product with a cannula, because at the exit, the product may enter the dermis layer.

Contraindications ⊘

- Pregnancy and lactation.
- Intolerance to components.
- Neuropsychiatric disorders, epilepsy.
- Wounds and inflammatory processes in the injection area.

Specification

Product name	Prime DR.LIPO+
Package unit	2 mL * 10 vials
Expiration date	36 months from date of manufacture. Check the box for the expiration date.
Storage	1~30℃
Ingredient	Water, Lecithin, Cyanocobalamin, Citric Acid, Sodium Citrate, Carnitine, EGF, FGF, Acetyl Hexapeptide-8, Palmitoyl Pentapeptide-4, Palmitoyl Tetrapeptide-7, Ascorbic Acid, Galloyl Pentapeptide-74, Nonapeptide-1, Glutathione, Adenosine, Arginine, Arbutin, 1,2-hexanediol.

BNC Global
Beauty and Culture
www.mybncglobal.com